

## PRODUCT SPECIFICATION

Product NumberAPrEST95453Gene Descriptionhydroxyacyl-thioester dehydratase type 2Corresponding AntibodiesAnti-HTD2 (HPA077228)DescriptionRecombinant protein fragment of Human HTD2Amino Acid SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: MFPLISSHHLWWGGLRRTVCLNLPVLTLQHFQHMHIKVGDRAELRRAFTQ TDVATFSELTGDVNPLHLNEDFAKHTKFGNTLVFusion TagN-terminal His <sub>6</sub> ABP (ABP = Albumin Binding Protein derived from Streptococcal Protein G)Expression Host <i>E. coli</i> PurificationIMAC chromatographyPredicted MW27 kDa including tagsUsageSuitable as control in WB and preadsorption assays using indicated corresponding antibodies.Purity>80% by SDS-PAGE and Coomassie blue stainingBufferPBS and 1M Urea, pH 7.4.Unit Size100 µlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.		
Gene Descriptionhydroxyacyl-thioester dehydratase type 2Corresponding AntibodiesAnti-HTD2 (HPA077228)DescriptionRecombinant protein fragment of Human HTD2Amino Acid SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: MFPLISSHHLWWGGLRRTVCLNLPVLTLQHFQHMHIKVGDRAELRRAFTQ TDVATFSELTGDVNPLHLNEDFAKHTKFGNTLVFusion TagN-terminal His <sub>6</sub> ABP (ABP = Albumin Binding Protein derived from Streptococcal Protein G)Expression Host <i>E. coli</i> PurificationIMAC chromatographyPredicted MW27 kDa including tagsUsageSuitable as control in WB and preadsorption assays using indicated corresponding antibodies.Purity>80% by SDS-PAGE and Coomassie blue stainingBufferPBS and 1M Urea, pH 7.4.Unit Size100 µlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Product Name	PrEST Antigen HTD2
Corresponding AntibodiesAnti-HTD2 (HPA077228)DescriptionRecombinant protein fragment of Human HTD2Amino Acid SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: MFPLISSHHLWWGGLRRTVCLNLPVLTLQHFQHMHIKVGDRAELRRAFTQ TDVATFSELTGDVNPLHLNEDFAKHTKFGNTLVFusion TagN-terminal His <sub>6</sub> ABP (ABP = Albumin Binding Protein derived from Streptococcal Protein G)Expression Host <i>E. coli</i> PurificationIMAC chromatographyPredicted MW27 kDa including tagsUsageSuitable as control in WB and preadsorption assays using indicated corresponding antibodies.Purity>80% by SDS-PAGE and Coomassie blue stainingBufferPBS and 1M Urea, pH 7.4.Unit Size100 µlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Product Number	APrEST95453
AntibodiesDescriptionRecombinant protein fragment of Human HTD2Amino Acid SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: MFPLISSHHLWWGGLRRTVCLNLPVLTLQHFQHMHIKVGDRAELRRAFTQ TDVATFSELTGDVNPLHLNEDFAKHTKFGNTLVFusion TagN-terminal His <sub>6</sub> ABP (ABP = Albumin Binding Protein derived from Streptococcal Protein G)Expression Host <i>E. coli</i> PurificationIMAC chromatographyPredicted MW27 kDa including tagsUsageSuitable as control in WB and preadsorption assays using indicated corresponding antibodies.Purity>80% by SDS-PAGE and Coomassie blue stainingBufferPBS and 1M Urea, pH 7.4.Unit Size100 μlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Gene Description	hydroxyacyl-thioester dehydratase type 2
Amino Acid SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: MFPLISSHHLWWGGLRRTVCLNLPVLTLQHFQHMHIKVGDRAELRRAFTQ TDVATFSELTGDVNPLHLNEDFAKHTKFGNTLVFusion TagN-terminal His <sub>6</sub> ABP (ABP = Albumin Binding Protein derived from Streptococcal Protein G)Expression Host <i>E. coli</i> PurificationIMAC chromatographyPredicted MW27 kDa including tagsUsageSuitable as control in WB and preadsorption assays using indicated corresponding antibodies.Purity>80% by SDS-PAGE and Coomassie blue stainingBufferPBS and 1M Urea, pH 7.4.Unit Size100 μlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Corresponding Antibodies	Anti-HTD2 (HPA077228)
MFPLISSHHLWWGGLRRTVCLNLPVLTLQHFQHMHIKVGDRAĚLRRAFTQ TDVATFSELTGDVNPLHLNEDFAKHTKFGNTLVFusion TagN-terminal His <sub>6</sub> ABP (ABP = Albumin Binding Protein derived from Streptococcal Protein G)Expression Host <i>E. coli</i> PurificationIMAC chromatographyPredicted MW27 kDa including tagsUsageSuitable as control in WB and preadsorption assays using indicated corresponding antibodies.Purity>80% by SDS-PAGE and Coomassie blue stainingBufferPBS and 1M Urea, pH 7.4.Unit Size100 μlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Description	Recombinant protein fragment of Human HTD2
G)Expression HostE. coliPurificationIMAC chromatographyPredicted MW27 kDa including tagsUsageSuitable as control in WB and preadsorption assays using indicated corresponding antibodies.Purity>80% by SDS-PAGE and Coomassie blue stainingBufferPBS and 1M Urea, pH 7.4.Unit Size100 μlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Amino Acid Sequence	MFPLISSHHLWWGGLRRTVCLNLPVLTLQHFQHMHIKVGDRAELRRAFTQ
PurificationIMAC chromatographyPredicted MW27 kDa including tagsUsageSuitable as control in WB and preadsorption assays using indicated corresponding antibodies.Purity>80% by SDS-PAGE and Coomassie blue stainingBufferPBS and 1M Urea, pH 7.4.Unit Size100 μlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Fusion Tag	
Predicted MW27 kDa including tagsUsageSuitable as control in WB and preadsorption assays using indicated corresponding antibodies.Purity>80% by SDS-PAGE and Coomassie blue stainingBufferPBS and 1M Urea, pH 7.4.Unit Size100 µlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Expression Host	E. coli
UsageSuitable as control in WB and preadsorption assays using indicated corresponding antibodies.Purity>80% by SDS-PAGE and Coomassie blue stainingBufferPBS and 1M Urea, pH 7.4.Unit Size100 µlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Purification	IMAC chromatography
antibodies.Purity>80% by SDS-PAGE and Coomassie blue stainingBufferPBS and 1M Urea, pH 7.4.Unit Size100 µlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Predicted MW	27 kDa including tags
BufferPBS and 1M Urea, pH 7.4.Unit Size100 µlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Usage	
Unit Size100 μlConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Purity	>80% by SDS-PAGE and Coomassie blue staining
ConcentrationLot dependentStorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Buffer	PBS and 1M Urea, pH 7.4.
StorageUpon delivery store at -20°C. Avoid repeated freeze/thaw cycles.NotesGently mix before use. Optimal concentrations and conditions for each application should	Unit Size	100 µl
Notes Gently mix before use. Optimal concentrations and conditions for each application should	Concentration	Lot dependent
	Storage	Upon delivery store at -20°C. Avoid repeated freeze/thaw cycles.
	Notes	

Product of Sweden. For research use only. Not intended for pharmaceutical development, diagnostic, therapeutic or any in vivo use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB.

Warranty: The products supplied by Atlas Antibodies are warranted to meet stated product specifications and to conform to label descriptions when used and stored properly. Unless otherwise stated, this warranty is limited to one year from date of sales for products used, handled and stored according to Atlas Antibodies AB's instructions. Atlas Antibodies AB's sole liability is limited to replacement of the product or refund of the purchase price. All products are supplied for research use only. They are not intended for medicinal, diagnostic or therapeutic use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB Rev. December 2012

IBAN

Bankgiro