

PRODUCT SPECIFICATION

Product Number HPA005456 Gene Description AT rich interactive domain 1A (SWI-like) Clonality Polyclonal Isotype IgG Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: PGLGNVAMGPRQHYPYGGPYDRVRTEPGIGPEGMMSTGAPQPNLMPSNPD SGMYSPSRYPPQQQQQQQQRHDSYGNQFSTQGTPSGSPFPSQQTTMYQQQ QQNYK Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Recommended Human HPC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml Characterization Data Available at atlasantibodies.com/products/HPA005456		
Gene Description AT rich interactive domain 1A (SWI-like) Clonality Polyclonal Isotype IgG Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: PGLGNVAMGPRQHYPYGGPYDRVRTEPGIGPEGNMSTGAPQPNLMPSNPD SGMYSPSRYPPQQQQQQQRHDSYGNQFSTQGTPSGSPFPSQQTTMYQQQ QQNYK Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Human Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml Characterization Data Available at atlasantibodies.com/products/HPA005456	Product Name	Anti-ARID1A
Clonality Polyclonal Isotype IgG Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: PGLGNVAMGPRQHYPYGGPYDRVRTEPGIGPEGNMSTGAPQPNLMPSNPD SGMYSPSRYPPQQQQQQQRHDSYGNQFSTQGTPSGSPFPSQQTTMYQQQ QQNYK Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Human Recommended Applications IHC (Immunohistochemistry)	Product Number	HPA005456
Host Rabbit	Gene Description	AT rich interactive domain 1A (SWI-like)
Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: PGLGNVAMGPRQHYPYGGPYDRVRTEPGIGPEGNMSTGAPQPNLMPSNPD SGMYSPSRYPPQQQQQQQRHDSYGNQFSTQGTPSGSPFPSQQTTMYQQQ QQNYK Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Human Recommended Applications HC (Immunohistochemistry)	Clonality	Polyclonal
Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: PGLGNVAMGPRQHYPYGGPYDRVRTEPGIGPEGNMSTGAPQPNLMPSNPD SGMYSPSRYPPQQQQQQQQRHDSYGNQFSTQGTPSGSPFPSQQTTMYQQQ QQNYK Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Recommended Human IHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml Characterization Data Available at atlasantibodies.com/products/HPA005456	Isotype	IgG
PGLGNVAMGPRQHYPYGGPYDRVRTEPGIGPEGNMSTGAPQPNLMPSNPD SGMYSPSRYPPQQQQQQQQRHDSYGNQFSTQGTPSGSPFPSQQTTMYQQQ QQNYK Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Recommended Human HC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml Characterization Data PGLGNVAMGPRQPNLMPSNPD SGMYSPSRYPPQQQQQQQQQRHDSYGNETSQPPNLMPSNPD SGMYSPSRYPPQQQQQQQQQQQRHDSYGNPSQPPNLMPSNPD SGMYSPSRYPPQQQQQQQQQQQQRHDSYGNQPPNLMPSNPD SGMYSPSRYPPQQQQQQQQQQQQQQQQQQQQQQQQQQQQQQQQ	Host	Rabbit
Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml Characterization Data Human Human Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml	Antigen Sequence	PGLGNVAMGPRQHYPYGGPYDRVRTEPGIGPEGNMSTGAPQPNLMPSNPD SGMYSPSRYPPQQQQQQQRHDSYGNQFSTQGTPSGSPFPSQQTTMYQQQ
Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml Characterization Data Available at atlasantibodies.com/products/HPA005456	Purification Method	Affinity purified using the PrEST antigen as affinity ligand
Applications - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml Characterization Data Available at atlasantibodies.com/products/HPA005456		Human
		- Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100
Doffers 400% above at DDO (at 1.7.0), 0.000% and its and its add at a process at its	Characterization Data	Available at atlasantibodies.com/products/HPA005456
Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.	Buffer	40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.
Concentration Lot dependent	Concentration	Lot dependent
Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.	Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C.
Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see atlasantibodies.com.	Notes	For protocols, additional product information, such as images and references, see

Product of Sweden. For research use only. Not intended for pharmaceutical development, diagnostic, therapeutic or any in vivo use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB.

Warranty: The products supplied by Atlas Antibodies are warranted to meet stated product specifications and to conform to label descriptions when used and stored properly. Unless otherwise stated, this warranty is limited to one year from date of sales for products used, handled and stored according to Atlas Antibodies AB's instructions. Atlas Antibodies AB's sole liability is limited to replacement of the product or refund of the purchase price. All products are supplied for research use only. They are not intended for medicinal, diagnostic or therapeutic use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB Rev. December 2012

Atlas Antibodies AB Phone +46(0)8 54 59 58 50 IBAN SE91 6000 0000 0004 6991 6761 Bankgiro 5469-1092