

PRODUCT SPECIFICATION

Product Number HPA009678 Gene Description GRAM domain containing 3 Clonality Polyclonal Isotype IgG Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: GKRESKLGSAHSEAENGVEEKKACRSPTAQSPTPSVEADSPDQKKIISL WSKSSFDGASLASDKNDCKTESKNDPKTERKKSSSSSQYKANMHFHKLFL SVPTEEPLKQSFTCALQKEILYQGKLFVSENWI Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Recommended Applications HIC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml Characterization Data Available at atlasantibodies.com/products/HPA009678 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see atlasantibodies.com.		
Gene Description Clonality Polyclonal Isotype IgG Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: GKRESKLGSAHSEAENGVEEKKKACRSPTAQSPTPSVEADSPDQKKIISL WSKSSFDGASLASDKNDCKTESKNDPKTERKKSSSSSQYKANMHFHKLFL SVPTEEPLKQSFTCALQKEILYQGKLFVSENWI Purification Method Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml Characterization Data Available at atlasantibodies.com/products/HPA009678 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Product Name	Anti-GRAMD3
Sotype IgG	Product Number	HPA009678
Isotype IgG Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: GKRESKLGSAHSEAENGVEEKKKACRSPTAQSPTPSVEADSPDQKKIISL WSKSSFDGASLASDKNDCKTESKNDPKTERKKSSSSSQYKANMHFHKLFL SVPTEEPLKQSFTCALQKEILYQGKLFVSENWI Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Human Recommended Applications Human Human HIC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml Characterization Data Available at atlasantibodies.com/products/HPA009678 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Gene Description	GRAM domain containing 3
Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: GKRESKLGSAHSEAENGVEEKKKACRSPTAQSPTPSVEADSPDQKKIISL WSKSSFDGASLASDKNDCKTESKNDPKTERKKSSSSSQYKANMHFHKLFL SVPTEEPLKQSFTCALQKEILYQGKLFVSENWI Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Recommended Human HC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml Characterization Data Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Clonality	Polyclonal
Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: GKRESKLGSAHSEAENGVEEKKKACRSPTAQSPTPSVEADSPDQKKIISL WSKSSFDGASLASDKNDCKTESKNDPKTERKKSSSSSQYKANMHFHKLFL SVPTEEPLKQSFTCALQKEILYQGKLFVSENWI Purification Method Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml Characterization Data Available at atlasantibodies.com/products/HPA009678 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Isotype	IgG
GKRESKLGSAHSEAENGVEEKKKACRSPTAQSPTPSVEADSPDQKKIISL WSKSSFDGASLASDKNDCKTESKNDPKTERKKSSSSQYKANMHFHKLFL SVPTEEPLKQSFTCALQKEILYQGKLFVSENWI Purification Method Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml Characterization Data Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Host	Rabbit
Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml Characterization Data Available at atlasantibodies.com/products/HPA009678 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Antigen Sequence	GKRESKLGSAHSEAENGVEEKKKÁCRSPTAQŠPTPSVEÁDSPĎQKKIÍSL WSKSSFDGASLASDKNDCKTESKNDPKTERKKSSSSSQYKANMHFHKLFL
Recommended Applications - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml Characterization Data Available at atlasantibodies.com/products/HPA009678 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Purification Method	Affinity purified using the PrEST antigen as affinity ligand
Applications - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml Characterization Data Available at atlasantibodies.com/products/HPA009678 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see		Human
Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see		- Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100
Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Characterization Data	Available at atlasantibodies.com/products/HPA009678
Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Buffer	40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.
Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Concentration	Lot dependent
be determined by the user. For protocols, additional product information, such as images and references, see	Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C.
	Notes	be determined by the user. For protocols, additional product information, such as images and references, see

Product of Sweden. For research use only. Not intended for pharmaceutical development, diagnostic, therapeutic or any in vivo use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB.

Warranty: The products supplied by Atlas Antibodies are warranted to meet stated product specifications and to conform to label descriptions when used and stored properly. Unless otherwise stated, this warranty is limited to one year from date of sales for products used, handled and stored according to Atlas Antibodies AB's instructions. Atlas Antibodies AB's sole liability is limited to replacement of the product or refund of the purchase price. All products are supplied for research use only. They are not intended for medicinal, diagnostic or therapeutic use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB Rev. December 2012

Atlas Antibodies AB Phone +46(0)8 54 59 58 50 IBAN SE91 6000 0000 0004 6991 6761 Bankgiro 5469-1092