

## PRODUCT SPECIFICATION

Atlas Antibodies AB

Phone

+46(0)8 54 59 58 50

IBAN

Product NumberHPA019123Gene DescriptionQKI, KH domain containing, RNA bindingClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: RAEIKLKRAVEEVKKLLVPAAEGEDSLKKMQLMELAILNGTYRDANIKSP ALAFSLAATAQAAPRIITGPAPVLPPAALRTPTPAGPTIMPLIRQIQTAV MPNGTPHPTAAIVPPGPEAGLIYTPYEYPYTLAPATSILEYPIEPSGVLPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityIHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml IICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization Data BufferAvailable at atlasantibodies.com/products/HPA019123Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependent	Product Name	Anti-OKI
Gene DescriptionQKI, KH domain containing, RNA bindingClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: RAEIKLKRAVEEVKKLLVPAAEGEDSLKKMQLMELAILNGTYRDANIKSP ALAFSLAATAQAAPRIITGPAPVLPPAALRTPTPAGPTIMPLIRQIQTAV MPNGTPHPTAAIVPPGPEAGLIYTPYEYPYTLAPATSILEYPIEPSGVLPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHuman, Mouse, RatRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization Data BufferAvailable at atlasantibodies.com/products/HPA019123Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Lot dependent		
ClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: RAETIKLKRAVEEVKKLLVPAAEGEDSLKKMQLMELATLNGTYRDANIKSP ALAFSLAATAQAAPRIITGPAPVLPPAALRTPTPAGPTIMPLIRQIQTAV MPNGTPHPTAAIVPPGPEAGLIYTPYEYPYTLAPATSILEYPIEPSGVLPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHuman, Mouse, RatRecommended ApplicationsHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA019123Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependent		
IsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: RAEIKLKRAVEEVKKLLVPAAEGEDSLKKMQLMELAILNGTYRDANIKSP ALAFSLAATAQAAPRIITGPAPVLPPAALRTPTPAGPTIMPLIRQIQTAV MPNGTPHPTAAIVPPGPEAGLIYTPYEYPYTLAPATSILEYPIEPSGVLPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHuman, Mouse, RatRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization Data BufferAvailable at atlasantibodies.com/products/HPA019123Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependent	Gene Description	QKI, KH domain containing, RNA binding
HostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: RAEIKLKRAVEEVKKLLVPAAEGEDSLKKMQLMELAILNGTYRDANIKSP ALAFSLAATAQAAPRIITGPAPVLPPAALRTPTPAGPTIMPLIRQIQTAV MPNGTPHPTAAIVPPGPEAGLIYTPYEYPYTLAPATSILEYPIEPSGVLPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHuman, Mouse, RatRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA019123Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependent	Clonality	Polyclonal
Antigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: RAEIKLKRAVEEVKKLLVPAAEGEDSLKKMQLMELAILNGTYRDANIKSP ALAFSLAATAQAAPRIITGPAPVLPPAALRTPTPAGPTIMPLIRQIQTAV MPNGTPHPTAAIVPPGPEAGLIYTPYEYPYTLAPATSILEYPIEPSGVLPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHuman, Mouse, RatRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization Data BufferAvailable at atlasantibodies.com/products/HPA019123Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependent	Isotype	lgG
RAEIKLKRAVEEVKKLLVPAAEGEDSLKKMQLMELAILNGTYRDANIKSP ALAFSLAATAQAAPRIITGPAPVLPPAALRTPTPAGPTIMPLIRQIQTAV MPNGTPHPTAAIVPPGPEAGLIYTPYEYPYTLAPATSILEYPIEPSGVLPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHuman, Mouse, RatRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization Data BufferAvailable at atlasantibodies.com/products/HPA019123Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependent	Host	Rabbit
Verified Species ReactivityHuman, Mouse, RatRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA019123Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependent	Antigen Sequence	RAEIKLKRAVEEVKKLLVPAAEGEDSLKKMQLMELAILNGTYRDANIKSP ALAFSLAATAQAAPRIITGPAPVLPPAALRTPTPAGPTIMPLIRQIQTAV
ReactivityRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 	Purification Method	Affinity purified using the PrEST antigen as affinity ligand
Applications- Antibody dilution: 1:500 - 1:1000 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA019123Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependent		Human, Mouse, Rat
Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependent		<ul> <li>Antibody dilution: 1:500 - 1:1000</li> <li>Retrieval method: HIER pH6</li> <li>WB (Western Blot)</li> <li>Working concentration: 0.04-0.4 μg/ml</li> <li>ICC-IF (Immunofluorescence)</li> <li>Fixation/Permeabilization: PFA/Triton X-100</li> </ul>
Concentration Lot dependent	Characterization Data	Available at atlasantibodies.com/products/HPA019123
	Buffer	40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.
Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.	Concentration	Lot dependent
	Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C.
Notes Gently mix before use. Optimal concentrations and conditions for each application shoul be determined by the user. For protocols, additional product information, such as images and references, see atlasantibodies.com.	Notes	For protocols, additional product information, such as images and references, see

Product of Sweden. For research use only. Not intended for pharmaceutical development, diagnostic, therapeutic or any in vivo use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB.

Warranty: The products supplied by Atlas Antibodies are warranted to meet stated product specifications and to conform to label descriptions when used and stored properly. Unless otherwise stated, this warranty is limited to one year from date of sales for products used, handled and stored according to Atlas Antibodies AB's instructions. Atlas Antibodies AB's sole liability is limited to replacement of the product or refund of the purchase price. All products are supplied for research use only. They are not intended for medicinal, diagnostic or therapeutic use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB Rev. December 2012

SE91 6000 0000 0004 6991 6761

Bankgiro

5469-1092