

PRODUCT SPECIFICATION

Product NumberHPA024679Gene Descriptiontonsoku-like, DNA repair proteinClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: WNRNDMGETLLHRACIEGQLRRVQDLVRQGHPLNPRDYCGWTPLHEACN YGHLEIVRFLLDHGAAVDDPGGQGCEGITPLHDALNCGHFEVAELLLERG ASVTLRTRKPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: 0:25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA024679Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. ConcentrationConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see		
Gene Descriptiontonsoku-like, DNA repair proteinClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: WNRRNDMGETLLHRACIEGQLRRVQDLVRQGHPLNPRDYCGWTPLHEACN YGHLETVRFLLDHGAAVDDPGQQCEGITPLHDALNCGHFEVAELLLERG ASVTLRTRKPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA024679Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Product Name	Anti-TONSL
ClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: WNRRNDMGETLLHRACIEGQLRRVQDLVRQGHPLNPRDYCGWTPLHEACN YGHLETVRFLLDHGAAVDDPGQQCEGITPLHDALNCGHFEVAELLLERG ASVTLRTRKPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA024679Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Product Number	HPA024679
IsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: WNRRNDMGETLLHRACIEGQLRRVQDLVRQGHPLNPRDYCGWTPLHEACN YGHLEIVRFLLDHGAAVDDPGGQGCEGITPLHDALNCGHFEVAELLLERG ASVTLRTRKPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA024679Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Gene Description	tonsoku-like, DNA repair protein
HostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: WNRRNDMGETLLHRACIEGQLRRVQDLVRQGHPLNPRDYCGWTPLHEACN YGHLEIVRFLLDHGAAVDDPGQGCEGITPLHDALNCGHFEVAELLLERG ASVTLRTRKPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization Data StorageAvailable at atlasantibodies.com/products/HPA024679Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Clonality	Polyclonal
Antigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: WNRNDMGETLLHRACIEGQLRRVQDLVRQGHPLNPRDYCGWTPLHEACN YGHLEIVRFLLDHGAAVDDPGGQGCEGITPLHDALNCGHFEVAELLLERG ASVTLRTRKPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization Data Store at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Isotype	lgG
WNRRNDMGETLLHRACIEGQLRRVQDLVRQGHPLNPRDÝCGWTPLHEÁCN YGHLEIVRFLLDHGAVDDPGGQGCEGITPLHDALNCGHFEVAELLLERG ASVTLRTRKPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA024679Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Host	Rabbit
Verified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA024679Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Antigen Sequence	WNRRNDMGETLLHRACIÉGQLRRVQDLVRQGHPLNPRDÝCGWTPLHEACN YGHLEIVRFLLDHGAAVDDPGGQGCEGITPLHDALNCGHFEVAELLLERG
ReactivityIHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) 	Purification Method	Affinity purified using the PrEST antigen as affinity ligand
Applications- Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA024679Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see		Human
Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see		- Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100
Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Characterization Data	Available at atlasantibodies.com/products/HPA024679
Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Buffer	40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.
Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Concentration	Lot dependent
be determined by the user. For protocols, additional product information, such as images and references, see	Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C.
atlasantibodies.com.	Notes	be determined by the user.

Product of Sweden. For research use only. Not intended for pharmaceutical development, diagnostic, therapeutic or any in vivo use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB.

Warranty: The products supplied by Atlas Antibodies are warranted to meet stated product specifications and to conform to label descriptions when used and stored properly. Unless otherwise stated, this warranty is limited to one year from date of sales for products used, handled and stored according to Atlas Antibodies AB's instructions. Atlas Antibodies AB's sole liability is limited to replacement of the product or refund of the purchase price. All products are supplied for research use only. They are not intended for medicinal, diagnostic or therapeutic use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB Rev. December 2012

Phone

Bankgiro