

## PRODUCT SPECIFICATION

Product Name Anti-CALCOCO1 Product Number HPA038313 Gene Description calcium binding and coiled-coil domain 1 Clonality Polyclonal Isotype IgG Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: EHTELMEQYKGISRSHGEITERDILSRQQGDHVARILELEDDIQTISEK VLTKEVELDRLRDTVKALTREQEKLLGQLKEVQADKEQSEAELQVAQQEN HHL  Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Recommended Applications Human  HC (Immunohistochemistry) - Antibody dilution: 1:200 - 1:500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml  Characterization Data Available at atlasantibodies.com/products/HPA038313  Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see atlasantibodies.com.		
Gene Description calcium binding and coiled-coil domain 1  Clonality Polyclonal  Isotype IgG  Host Rabbit  Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: EHTELMEQYKGISRSHGEITEERDILSRQQGDHVARILELEDDIQTISEK VLTKEVELDRLRDTVKALTREQEKLLGQLKEVQADKEQSEAELQVAQQEN HHL  Purification Method Affinity purified using the PrEST antigen as affinity ligand  Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:200 - 1:500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml  Characterization Data Available at atlasantibodies.com/products/HPA038313  Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Lot dependent  Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Product Name	Anti-CALCOCO1
Clonality   Polyclonal   IgG	Product Number	HPA038313
Isotype	Gene Description	calcium binding and coiled-coil domain 1
Host       Rabbit         Antigen Sequence       Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: EHTELMEQYKGISRSHGEITEERDILSRQQGDHVARILELEDDIQTISEK VLTKEVELDRLRDTVKALTREQEKLLGQLKEVQADKEQSEAELQVAQQEN HHL         Purification Method       Affinity purified using the PrEST antigen as affinity ligand         Verified Species Reactivity       Human         Recommended Applications       IHC (Immunohistochemistry)         - Antibody dilution: 1:200 - 1:500       - Retrieval method: HIER pH6         WB (Western Blot)       - Working concentration: 0.04-0.4 μg/ml         ICC-IF (Immunofluorescence)       - Fixation/Permeabilization: PFA/Triton X-100         - Working concentration: 0.25-2 μg/ml         Characterization Data       Available at atlasantibodies.com/products/HPA038313         Buffer       40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.         Concentration       Lot dependent         Store at +4°C for short term storage. Long time storage is recommended at -20°C.         Notes       Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.         For protocools, additional product information, such as images and references, see	Clonality	Polyclonal
Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: EHTELMEQYKGISRSHGEITERDILSRQQGDHVARILELEDDIQTISEK VLTKEVELDRLRDTVKALTREQEKLLGQLKEVQADKEQSEAELQVAQQEN HHL  Purification Method Affinity purified using the PrEST antigen as affinity ligand  Verified Species Reactivity  Recommended Applications  IHC (Immunohistochemistry) - Antibody dilution: 1:200 - 1:500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml  Characterization Data Available at atlasantibodies.com/products/HPA038313  Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Isotype	IgG
EHTELMEQYKGISRSHGEITEERDILSRQQGDHVARILELEDDIQTISEK VLTKEVELDRLRDTVKALTREQEKLLGQLKEVQADKEQSEAELQVAQQEN HHL  Purification Method Affinity purified using the PrEST antigen as affinity ligand  Verified Species Reactivity  Recommended Human  HC (Immunohistochemistry) - Antibody dilution: 1:200 - 1:500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml  Characterization Data Available at atlasantibodies.com/products/HPA038313  Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Host	Rabbit
Verified Species Reactivity       Human         Recommended Applications       IHC (Immunohistochemistry)         Applications       - Antibody dilution: 1:200 - 1:500         - Retrieval method: HIER pH6 WB (Western Blot)       - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence)         - Fixation/Permeabilization: PFA/Triton X-100       - Working concentration: 0.25-2 μg/ml         Characterization Data       Available at atlasantibodies.com/products/HPA038313         Buffer       40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.         Concentration       Lot dependent         Storage       Store at +4°C for short term storage. Long time storage is recommended at -20°C.         Notes       Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.         For protocols, additional product information, such as images and references, see	Antigen Sequence	EHTELMEQYKGISRSHGEITEERDILSRQQGDHVARILELEDDIQTISEK VLTKEVELDRLRDTVKALTREQEKLLGQLKEVQADKEQSEAELQVAQQEN
Recommended Applications  - Antibody dilution: 1:200 - 1:500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml  Characterization Data Available at atlasantibodies.com/products/HPA038313  Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Purification Method	Affinity purified using the PrEST antigen as affinity ligand
Applications  - Antibody dilution: 1:200 - 1:500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml  Characterization Data Available at atlasantibodies.com/products/HPA038313  Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Lot dependent  Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see		Human
Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Lot dependent  Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.  For protocols, additional product information, such as images and references, see		- Antibody dilution: 1:200 - 1:500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100
Concentration  Storage  Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes  Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.  For protocols, additional product information, such as images and references, see	Characterization Data	Available at atlasantibodies.com/products/HPA038313
Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.  For protocols, additional product information, such as images and references, see	Buffer	40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.
Notes  Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.  For protocols, additional product information, such as images and references, see	Concentration	Lot dependent
be determined by the user.  For protocols, additional product information, such as images and references, see	Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C.
	Notes	be determined by the user. For protocols, additional product information, such as images and references, see

Product of Sweden. For research use only. Not intended for pharmaceutical development, diagnostic, therapeutic or any in vivo use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB.

Warranty: The products supplied by Atlas Antibodies are warranted to meet stated product specifications and to conform to label descriptions when used and stored properly. Unless otherwise stated, this warranty is limited to one year from date of sales for products used, handled and stored according to Atlas Antibodies AB's instructions. Atlas Antibodies AB's sole liability is limited to replacement of the product or refund of the purchase price. All products are supplied for research use only. They are not intended for medicinal, diagnostic or therapeutic use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB Rev. December 2012

Atlas Antibodies AB Phone +46(0)8 54 59 58 50 IBAN SE91 6000 0000 0004 6991 6761 Bankgiro 5469-1092