

PRODUCT SPECIFICATION

| Product NameAnti-FTLProduct NumberHPA041602Gene Descriptionferritin, light polypeptideClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: LFQDIKKPAEDEWGKTPDAMKAAMALEKKLNQALLDLHALGSARTDPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended Applications- Antibody dilution: 1:1000 - 1:2500 - Retrieval method: HIER pH6 WB (Western Blot) - Vorking concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization Data StorageAvailable at atlasantibodies.com/products/HPA041602Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.Concentration StorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see atlasantibodies.com. | | |
|--|-----------------------|--|
| Gene Descriptionferritin, light polypeptideClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: LFQDIKKPAEDEWGKTPDAMKAAMALEKKLNQALLDLHALGSARTDPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommendedHLC (Immunohistochemistry) - Antibody dilution: 1:1000 - 1:2500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofilourescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA041602Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | Product Name | Anti-FTL |
| ClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: LFQDIKKPAEDEWGKTPDAMKAAMALEKKLNQALLDLHALGSARTDPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:1000 - 1:2500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA041602Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | Product Number | HPA041602 |
| IsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: LFQDIKKPAEDEWGKTPDAMKAAMALEKKLNQALLDLHALGSARTDPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:1000 - 1:2500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA041602Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | Gene Description | ferritin, light polypeptide |
| HostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: LFQDIKKPAEDEWGKTPDAMKAAMALEKKLNQALLDLHALGSARTDPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:1000 - 1:2500 - Retrieval method: HIER PH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA041602Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | Clonality | Polyclonal |
| Antigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: LFQDIKKPAEDEWGKTPDAMKAAMALEKKLNQALLDLHALGSARTDPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsHC (Immunohistochemistry) - Antibody dilution: 1:1000 - 1:2500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA041602Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | Isotype | lgG |
| LFQDIKKPAEDEWGKTPDAMKAAMALEKKLNQALLDLHALGSARTDPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:1000 - 1:2500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA041602Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | Host | Rabbit |
| Verified Species Reactivity Human Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:1000 - 1:2500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml Characterization Data Available at atlasantibodies.com/products/HPA041602 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | Antigen Sequence | |
| ReactivityRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:1000 - 1:2500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 µg/ml ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA041602Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | Purification Method | Affinity purified using the PrEST antigen as affinity ligand |
| Applications- Antibody dilution: 1:1000 - 1:2500 - Retrieval method: HIER pH6 WB (Western Blot) - Working concentration: 0.04-0.4 μg/ml | | Human |
| Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | | Antibody dilution: 1:1000 - 1:2500 Retrieval method: HIER pH6 WB (Western Blot) Working concentration: 0.04-0.4 μg/ml ICC-IF (Immunofluorescence) Fixation/Permeabilization: PFA/Triton X-100 |
| Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | Characterization Data | Available at atlasantibodies.com/products/HPA041602 |
| Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | Buffer | 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. |
| Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see | Concentration | Lot dependent |
| be determined by the user. For protocols, additional product information, such as images and references, see | Storage | Store at +4°C for short term storage. Long time storage is recommended at -20°C. |
| | Notes | be determined by the user. For protocols, additional product information, such as images and references, see |

Product of Sweden. For research use only. Not intended for pharmaceutical development, diagnostic, therapeutic or any in vivo use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB.

Warranty: The products supplied by Atlas Antibodies are warranted to meet stated product specifications and to conform to label descriptions when used and stored properly. Unless otherwise stated, this warranty is limited to one year from date of sales for products used, handled and stored according to Atlas Antibodies AB's instructions. Atlas Antibodies AB's sole liability is limited to replacement of the product or refund of the purchase price. All products are supplied for research use only. They are not intended for medicinal, diagnostic or therapeutic use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB Rev. December 2012

Phone

Bankgiro