

## PRODUCT SPECIFICATION

Product NameAnti-PHC2Product NumberHPA047403Gene Descriptionpolyhomeotic homolog 2 (Drosophila)ClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: PQLASVSPSVALQPSSEAHAMPLGPVTPALPLQCPTANLHKPGGSQQCHP PTPDTGPQNGHPEGVPHTPQRRFQHTSAVILQLQPASPVPQQCVVPDDWKE VAPGEKSVPETRSGPSPHQQAIVTAMPPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsHC (Immunohistochemistry) - Antibody dilution: 1:20 - 1:50 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: DFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA047403Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see atlasantibodies.com.		
Gene Descriptionpolyhomeotic homolog 2 (Drosophila)ClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: PQLASVSPSVALQPSSEAHAMPLGPVTPALPLQCPTANLHKPGGSQQCHP PTPDTGPQNGHPEGVPHTPQRFQHTSAVILQLQPASPVPQQCVPDDWKE VAPGEKSVPETRSGPSPHQQAIVTAMPPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended Applications- Antibody dilution: 1:20 - 1:50 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization Data Available at atlasantibodies.com/products/HPA047403Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.Concentration StorageLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Product Name	Anti-PHC2
ClonalityPolyclonalIsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: PQLASVSPSVALQPSSEAHAMPLGPVTPALPLQCPTANLHKPGGSQQCHP PTPDTGPQNGHPEGVPHTPQRRFQHTSAVILQLQPASPVPQQCVPDDWKE VAPGEKSVPETRSGPSPHQQAIVTAMPPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsHC (Immunohistochemistry) - Antibody dilution: 1:20 - 1:50 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA047403Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Product Number	HPA047403
IsotypeIgGHostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: PQLASVSPSVALQPSSEAHAMPLGPVTPALPLQCPTANLHKPGGSQQCHP PTPDTGPQNGHPEGVPHTPQRRFQHTSAVILQLQPASPVPQQCVPDDWKE VAPGEKSVPETRSGPSPHQQAIVTAMPPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:20 - 1:50 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA047403Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. - For protocols, additional product information, such as images and references, see	Gene Description	polyhomeotic homolog 2 (Drosophila)
HostRabbitAntigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: PQLASVSPSVALQPSSEAHAMPLGPVTPALPLQCPTANLHKPGGSQQCHP PTPDTGPQNGHPEGVPHTPQRRFQHTSAVILQLQPASPVPQQCVPDDWKE VAPGEKSVPETRSGPSPHQQAIVTAMPPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:20 - 1:50 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA047403Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Clonality	Polyclonal
Antigen SequenceRecombinant Protein Epitope Signature Tag (PrEST) antigen sequence: PQLASVSPSVALQPSSEAHAMPLGPVTPALPLQCPTANLHKPGGSQQCHP PTPDTGPQNGHPEGVPHTPQRRFQHTSAVILQLQPASPVPQQCVPDDWKE VAPGEKSVPETRSGPSPHQQAIVTAMPPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:20 - 1:50 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: DFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization Data SufferAvailable at atlasantibodies.com/products/HPA047403Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.Concentration StorageLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Isotype	lgG
PQLASVSPSVALQPSSEAHAMPLGPVTPALPLQCPTANLHKPGGSQQCHP PTPDTGPQNGHPEGVPHTPQRRFQHTSAVILQLQPASPVPQQCVPDDWKE VAPGEKSVPETRSGPSPHQQAIVTAMPPurification MethodAffinity purified using the PrEST antigen as affinity ligandVerified Species ReactivityHumanRecommended ApplicationsHLC (Immunohistochemistry) - Antibody dilution: 1:20 - 1:50 - Retrieval method: HIER pH6 ICCC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA047403Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Host	Rabbit
Verified Species ReactivityHumanRecommended ApplicationsIHC (Immunohistochemistry) - Antibody dilution: 1:20 - 1:50 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA047403Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Antigen Sequence	PQLASVSPSVALQPSSEAHAMPLGPVTPALPLQCPTANLHKPGGSQQCHP PTPDTGPQNGHPEGVPHTPQRRFQHTSAVILQLQPASPVPQQCVPDDWKE
ReactivityIHC (Immunohistochemistry) - Antibody dilution: 1:20 - 1:50 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) 	Purification Method	Affinity purified using the PrEST antigen as affinity ligand
Applications- Antibody dilution: 1:20 - 1:50 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/mlCharacterization DataAvailable at atlasantibodies.com/products/HPA047403Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see		Human
Buffer40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see		- Antibody dilution: 1:20 - 1:50 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100
ConcentrationLot dependentStorageStore at +4°C for short term storage. Long time storage is recommended at -20°C.NotesGently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Characterization Data	Available at atlasantibodies.com/products/HPA047403
Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.   Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.   For protocols, additional product information, such as images and references, see	Buffer	40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.
Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Concentration	Lot dependent
be determined by the user. For protocols, additional product information, such as images and references, see	Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C.
	Notes	be determined by the user. For protocols, additional product information, such as images and references, see

Product of Sweden. For research use only. Not intended for pharmaceutical development, diagnostic, therapeutic or any in vivo use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB.

Warranty: The products supplied by Atlas Antibodies are warranted to meet stated product specifications and to conform to label descriptions when used and stored properly. Unless otherwise stated, this warranty is limited to one year from date of sales for products used, handled and stored according to Atlas Antibodies AB's instructions. Atlas Antibodies AB's sole liability is limited to replacement of the product or refund of the purchase price. All products are supplied for research use only. They are not intended for medicinal, diagnostic or therapeutic use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB Rev. December 2012

Phone

Bankgiro