

PRODUCT SPECIFICATION

Product Number Gene Description Zinc finger protein 426 Clonality Polyclonal Isotype IgG Host Antigen Sequence EKTPAGRTVADCLTDCYQDSVTFDDVAVDFTQEEWTLLDSTQRSLYSDVM LENYKNLATVGGQIIKPSLISWLEQEESRTVQGGVLQGWEMRLETQWSIL QQDFLRGQTSIGIQLEGKHNGRELC Purification Method Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml ChIP (Chromatin Immunoprecipitation) - Working concentration: 1-10µg per reaction - *Please note not all antibody batches are validated for use in ChIP applications Characterization Data Available at atlasantibodies.com/products/HPA029342 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentration, such as images and references, see	Due do et Nieue	A. F. ZAIFACO
Gene Description zinc finger protein 426 Clonality Polyclonal Isotype IgG Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: EKTPAGRIVADCLTDCYQDSVTFDDVAVDFTQEEWTLLDSTQRSLYSDVM LENYKNLATVGGQTIKPSLTSWLEQEESRTVQGGVLQGWEMRLETQWSIL QQDFLRGQTSIGIQLEGKHNGRELC Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Recommended Applications - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml ChIP (Chromatin Immunoprecipitation) - Working concentration: 1-10µg per reaction - *Please note not all antibody batches are validated for use in ChIP applications Characterization Data Available at atlasantibodies.com/products/HPA029342 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Product Name	Anti-ZNF426
Clonality Polyclonal Isotype IgG Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: EKTPAGRIVADCLTDCYQDSVTFDDVAVDFTQEWTLLDSTQRSLYSDVM LENYKNLATVGQTIKPSLISWLEQEESRTVQGGVLQGWEMRLETQWSIL QQDFLRGQTSIGIQLEGKHNGRELC Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml ChIP (Chromatin Immunoprecipitation) - Working concentration: 1-10µg per reaction - *Please note not all antibody batches are validated for use in ChIP applications Characterization Data Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Product Number	HPA029342
Isotype	Gene Description	zinc finger protein 426
Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: EKTPAGRIVADCLTDCYQDSVTFDDVAVDFTQEEWTLLDSTQRSLYSDVM LENYKNLATVGGQTIKPSLISWLEQEESRTVQGGVLQGWEMRLETQWSIL QQDFLRGQTSIGIQLEGKHNGRELC Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Recommended Applications HC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml ChIP (Chromatin Immunoprecipitation) - Working concentration: 1-10µg per reaction - *Please note not all antibody batches are validated for use in ChIP applications Characterization Data Available at atlasantibodies.com/products/HPA029342 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Clonality	Polyclonal
Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: EKTPAGRIVADCLTDCYQDSVTFDDVAVDFTQEEWTLLDSTQRSLYSDVM LENYKNLATVGGQIIKPSLISWLEQEESRTVQGGVLQGWEMRLETQWSIL QQDFLRGQTSIGIQLEGKHNGRELC Purification Method Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml ChIP (Chromatin Immunoprecipitation) - Working concentration: 1-10µg per reaction - *Please note not all antibody batches are validated for use in ChIP applications Characterization Data Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Isotype	lgG
EKTPAGRIVADCLTDCYQDSVTFDDVAVDFTQEEWTLLDSTQRSLYSDVM LENYKNLATVGGQIIKPSLISWLEQEESRTVQGGVLQGWEMRLETQWSIL QQDFLRGQTSIGIQLEGKHNGRELC Purification Method Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml ChIP (Chromatin Immunoprecipitation) - Working concentration: 1-10µg per reaction - *Please note not all antibody batches are validated for use in ChIP applications Characterization Data Buffer Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes EKTPAGRIVADCLTDCYQDSVTFDDVAVDFTQEEWTLLDSTQRSLYSDVM LENYKNLETQUSTLEQUSTLA Human Hum	Host	Rabbit
Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml ChIP (Chromatin Immunoprecipitation) - Working concentration: 1-10μg per reaction - *Please note not all antibody batches are validated for use in ChIP applications Characterization Data Available at atlasantibodies.com/products/HPA029342 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Antigen Sequence	EKTPAGRIVADCLTDCYQDSVTFDDVAVDFTQEEWTLLDSTQRSLYSDVM LENYKNLATVGGQIIKPSLISWLEQEESRTVQGGVLQGWEMRLETQWSIL
Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml ChIP (Chromatin Immunoprecipitation) - Working concentration: 1-10µg per reaction - *Please note not all antibody batches are validated for use in ChIP applications Characterization Data Available at atlasantibodies.com/products/HPA029342 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Purification Method	Affinity purified using the PrEST antigen as affinity ligand
Applications - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml ChIP (Chromatin Immunoprecipitation) - Working concentration: 1-10μg per reaction - *Please note not all antibody batches are validated for use in ChIP applications Characterization Data Available at atlasantibodies.com/products/HPA029342 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see		Human
Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Lot dependent Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see		- Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml ChIP (Chromatin Immunoprecipitation) - Working concentration: 1-10µg per reaction
Concentration Lot dependent Storage Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Characterization Data	Available at atlasantibodies.com/products/HPA029342
Storage Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Buffer	40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.
Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Concentration	Lot dependent
be determined by the user. For protocols, additional product information, such as images and references, see	Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C.
atiasantibodies.com.	Notes	be determined by the user.

Product of Sweden. For research use only. Not intended for pharmaceutical development, diagnostic, therapeutic or any in vivo use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB.

Warranty: The products supplied by Atlas Antibodies are warranted to meet stated product specifications and to conform to label descriptions when used and stored properly. Unless otherwise stated, this warranty is limited to one year from date of sales for products used, handled and stored according to Atlas Antibodies AB's instructions. Atlas Antibodies AB's sole liability is limited to replacement of the product or refund of the purchase price. All products are supplied for research use only. They are not intended for medicinal, diagnostic or therapeutic use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB Rev. December 2012

Atlas Antibodies AB Phone +46(0)8 54 59 58 50 IBAN SE91 6000 0000 0004 6991 6761 Bankgiro 5469-1092