

## PRODUCT SPECIFICATION

Product Number HPA064047  Gene Description solute carrier family 35 (UDP-GlcNAc/UDP-glucose transporter), member D2  Clonality Polyclonal  Isotype IgG  Host Rabbit  Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence:		
Gene Description solute carrier family 35 (UDP-GlcNAc/UDP-glucose transporter), member D2  Clonality Polyclonal Isotype IgG Host Rabbit  Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence:	Product Name	Anti-SLC35D2
Clonality Polyclonal Isotype IgG Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: YSFLTLSSQLKPKPVGEENICLD Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml  Characterization Data Available at atlasantibodies.com/products/HPA064047  Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Product Number	HPA064047
Isotype IgG Host Rabbit Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: YSFLTLSSQLKPKPVGEENICLD Purification Method Affinity purified using the PrEST antigen as affinity ligand Verified Species Reactivity Recommended Applications IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunohiuorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml  Characterization Data Available at atlasantibodies.com/products/HPA064047 Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative. Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C. Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Gene Description	solute carrier family 35 (UDP-GlcNAc/UDP-glucose transporter), member D2
Host       Rabbit         Antigen Sequence       Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: YSFLTLSSQLKPKPVGEENTCLD         Purification Method       Affinity purified using the PrEST antigen as affinity ligand         Verified Species Reactivity       Human         Recommended Applications       IHC (Immunohistochemistry) <ul> <li>- Antibody dilution: 1:50 - 1:200</li> <li>- Retrieval method: HIER pH6</li> <li>ICC-IF (Immunofluorescence)</li> <li>- Fixation/Permeabilization: PFA/Triton X-100</li> <li>- Working concentration: 0.25-2 μg/ml</li> </ul> Characterization Data     Available at atlasantibodies.com/products/HPA064047         Buffer       40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.         Concentration       Lot dependent         Storage       Store at +4°C for short term storage. Long time storage is recommended at -20°C.         Notes       Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.         For protocols, additional product information, such as images and references, see	Clonality	Polyclonal
Antigen Sequence Recombinant Protein Epitope Signature Tag (PrEST) antigen sequence: YSFLTLSSQLKPKPVGEENICLD  Purification Method Verified Species Reactivity Recommended Applications  IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 µg/ml  Characterization Data Available at atlasantibodies.com/products/HPA064047  Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Isotype	IgG
Purification Method Affinity purified using the PrEST antigen as affinity ligand  Verified Species Reactivity  Recommended Applications IHC (Immunohistochemistry)  Applications - Antibody dilution: 1:50 - 1:200  - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence)  - Fixation/Permeabilization: PFA/Triton X-100  - Working concentration: 0.25-2 µg/ml  Characterization Data Available at atlasantibodies.com/products/HPA064047  Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Lot dependent  Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.  For protocols, additional product information, such as images and references, see	Host	Rabbit
Verified Species Reactivity  Recommended Applications  IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml  Characterization Data Available at atlasantibodies.com/products/HPA064047  Buffer  40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes  Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Antigen Sequence	
Reactivity  Recommended Applications  IHC (Immunohistochemistry) - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml  Characterization Data Available at atlasantibodies.com/products/HPA064047  Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Lot dependent  Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	Purification Method	Affinity purified using the PrEST antigen as affinity ligand
Applications  - Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100 - Working concentration: 0.25-2 μg/ml  Characterization Data Available at atlasantibodies.com/products/HPA064047  Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user. For protocols, additional product information, such as images and references, see	•	Human
Buffer 40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.  Concentration Lot dependent  Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.  For protocols, additional product information, such as images and references, see		- Antibody dilution: 1:50 - 1:200 - Retrieval method: HIER pH6 ICC-IF (Immunofluorescence) - Fixation/Permeabilization: PFA/Triton X-100
Concentration  Lot dependent  Storage  Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes  Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.  For protocols, additional product information, such as images and references, see	Characterization Data	Available at atlasantibodies.com/products/HPA064047
Storage Store at +4°C for short term storage. Long time storage is recommended at -20°C.  Notes Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.  For protocols, additional product information, such as images and references, see	Buffer	40% glycerol and PBS (pH 7.2). 0.02% sodium azide is added as preservative.
Notes  Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.  For protocols, additional product information, such as images and references, see	Concentration	Lot dependent
be determined by the user.  For protocols, additional product information, such as images and references, see	Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C.
	Notes	For protocols, additional product information, such as images and references, see

Product of Sweden. For research use only. Not intended for pharmaceutical development, diagnostic, therapeutic or any in vivo use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB.

Warranty: The products supplied by Atlas Antibodies are warranted to meet stated product specifications and to conform to label descriptions when used and stored properly. Unless otherwise stated, this warranty is limited to one year from date of sales for products used, handled and stored according to Atlas Antibodies AB's instructions. Atlas Antibodies AB's sole liability is limited to replacement of the product or refund of the purchase price. All products are supplied for research use only. They are not intended for medicinal, diagnostic or therapeutic use. No products from Atlas Antibodies may be resold, modified for resale or used to manufacture commercial products without prior written approval from Atlas Antibodies AB Rev. December 2012

Atlas Antibodies AB Phone +46(0)8 54 59 58 50 IBAN SE91 6000 0000 0004 6991 6761 Bankgiro 5469-1092